

# Carbon chemistry in Early Earth's magma ocean

Natalia V. Solomatova<sup>1</sup>, Razvan Caracas<sup>1</sup> and Craig E. Manning<sup>2</sup>

<sup>1</sup>CNRS, École Normale Supérieure de Lyon, Université Claude Bernard Lyon 1, Laboratoire de Géologie de Lyon, CNRS UMR 5276, Lyon, France.

<sup>2</sup>Department of Earth, Planetary and Space Sciences, University of California, Los Angeles, USA.

The amount of carbon stored in closed hidden reservoirs, particularly the lower mantle and iron core, is currently uncertain, and estimates of the carbon flux between the surface and mantle are model-dependent. To improve our understanding of Earth's global carbon cycle, it is critical to characterize the forms in which carbon existed in the proto-lunar disk after the Giant Impact and in the molten early Earth. Here we implement first-principles molecular dynamics to study the evolution of carbon species as a function of pressure (0-140 GPa) and temperature (3000-5000 K) in pyrolite melts with 3.4-6.5 wt.% CO and 5.2-9.8 wt.% CO<sub>2</sub>, capturing the range of carbon concentrations in carbonaceous chondrites. Carbon displays a variety of coordination environments, which are affected by the pressure, temperature and oxidation state of the melt. At densities of the proto-lunar disk, we observe the formation of bubbles within the silicate melt that contain CO and CO<sub>2</sub> species. We characterize the speciation of all volatile species and quantify the amount of carbon in the bubbles compared to the silicate melt to provide insight into the fate of carbon after the Giant Impact. At ambient pressure, carbon exists in the melt exclusively as a mixture of CO, CO<sub>2</sub> and CO<sub>3</sub> (unpolymerized carbon) while at higher pressures, the abundance of CO<sub>2</sub> and CO<sub>3</sub> species decreases at the expense of CO<sub>4</sub> species and complex oxo-carbon polymers (C<sub>x</sub>O<sub>y</sub>). With increasing pressure, decreasing temperature and decreasing oxygen fugacity, carbon becomes increasingly polymerized, forming multiple carbon-carbon bonds, direct carbon-silicon bonds and complex iron-carbon polymers. The presence of iron-carbon clusters suggests that upon segregation, iron-rich metal may partition a significant fraction of carbon from the silicate liquid, leading to carbon transport into the Earth's core.

Acknowledgements: This research was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement n°681818 – IMPACT). The ab initio simulations were performed on the GENCI supercomputers, under eDARI/CINES grants x106368.